ATTY. DOCKET NUMBER: REF/DEHN

Attorney for Applicant

Registration Number: 26,382

# ECOMMISSIONER OF PATENTS AND TRADEMARKS



					04	1/21/2.
Transmitted herewith for	filing under 35 USC 11	1 and 37 CFR 1.53 is the	e 🗆 Design 🛭	Utility	patent applicat	ion of:
Inventor: Trygve GULBRANDSEN et al.						
ENTITLED: Proc	cess for the Preparation	of Contrast Agents				
Enclosed are:						
$\boxtimes 12$ pages of	$\boxtimes$ 12 pages of written description, claims and abstract.					
$\Box$ 0 sheet(s) of	□ 0 sheet(s) of drawings.					
An assignment of the invention to NYCOMED IMAGING AS						
X Executed dec	laration of the inventors	•				
☐ A certified co	opy of a	application. Priority	y is claimed if no	t already	y of record.	
☐ A verified sta	atement to establish smal	l entity status under 37 (	CFR 1.9 and 37 (	CFR 1.2	7.	
N Preliminary a	mendment.					
and the second s						
The second secon						
The filing fee has been c	alculated as shown below	v:				
ITEM A	S FILED	# EXTRA	SMALL ENT	гіту	FULL I	TEE
Basic Fee				\$385. \$160.	X Utility  Design	\$770. \$320.
Total Claims	12 - 20 =	<b>0</b> 1	× \$ 11 =		× \$ 22 =	0
Independent Claims	2 - 3 =	0 2	× \$ 40 =		× \$ 80 =	0
☐ Multiple Dependent Claims in Proper Form Presented			+ \$130 =		+ \$260 =	0
TOTAL			\$		\$	770
	<sup>1</sup> If less than 20	filed, enter 0. <sup>2</sup> If less than	3 filed, enter 0.			
□ Place charge my Do	nosit Assount Number	02 0200 in the amount a	£ ¢	4		- C ( 1
☐ Please charge my De assignment recording		te copy of this paper is en		то	cover the filin	g ree (and
	,, //	or this paper is the	acios <b>cu</b> ,			
X A check in the amount	nt of \$810.00 to cover	r the filing fee and assign	ment recording f	èe is en	closed.	
The Commissioner is	hereby authorized to cha-	rge any additional fees ass	sociated with this	commu	nication inclu	dina natant
The Commissioner is hereby authorized to charge any additional fees associated with this communication, including patent application filing fees and processing fees under 37 CFR 1.16 and 37 CFR 1.17 or credit any overpayment to Deposit						
Account Number 02-0200. A duplicate copy of this paper is enclosed.						
BACON & THOMAS Respectfully submitted,						
625 SLATERS LANE - FOURTH FLOOR						
ALEXANDRIA, VIRGII	NIA 22314					
(703) 683-0500			Riden.	05	Fields	<i>!</i> 
DATE:April 21, 1997			Richard E. Fi		, -cue	<u>r</u>

APPLIC.TML (WP51-7OcT96)

770.00 101 H

- 1 -

64974/002.594

### Process for the preparation of contrast agents

#### Field of the Invention

The invention relates to a process for the preparation of iodinated X-ray contrast agents, in particular non-ionic N-alkylated-acylamino-triiodophenyl compounds such as iohexol, iodixanol, iopentol and ioxilan.

#### Background of the Invention

Non-ionic iodinated X-ray contrast agents have achieved great commercial success over the past twenty years and accordingly improving the efficiency of their manufacture is of great importance.

The manufacture of non-ionic contrast agents includes production of the chemical drug substance (primary production), followed by formulation to drug product (secondary production). The drug substance is usually made in a multistep chemical synthesis, and is thoroughly purified before formulation. As with any commercial drug production it is important to optimize yield, process time and demand for expensive equipment. All these parameters depend both on the chemical reaction conditions and the work-up between each step. The number of steps in the overall synthesis will of course be of great importance and if the work-up between individual process steps could be omitted significant improvement in efficiency can be obtained, provided that sufficient quality and yield are maintained for the final product.

Omitting the work-up between two process steps means that the second step will be performed in the crude

reaction solution resulting from the previous step. Potential major advantages includes:

- higher yield (through avoidance of loss of the intermediate in the work-up);
- elimination of isolation, purification, drying and analysis of the intermediate;
- significant reduction in equipment demand (eg. reactors, filters, driers, etc.); and
- significant reduction of the process time.

Despite these potential advantages, in practice intermediates are usually worked up between successive process steps. The reasons for this often include the facts that impurities may carry through the process and that the optimum solvent for the first reaction is usually not the same as for the second reaction.

The present multistage preparation of certain non-ionic X-ray contrast agents requires successive acylation and N-alkylation reactions. Thus for example in the production of iohexol (as described in SE-7706792-4 and by Gulbrandsen in Kjemi No. 6/90, pages 6-8), 5-amino-N,N'-bis(2,3-dihydroxypropyl)-2,4,6-triiodophthalamide (hereinafter 5-Amine) is acetylated to produce 5-acetamido-N,N'-bis(2,3-dihydroxypropyl)-2,4,6-triiodophthalamide (hereinafter 5-Acetamide) which is then N-alkylated to produce 5-[N-(2,3-dihydroxypropyl)-acetamido]-N,N'-bis(2,3-dihydroxypropyl)-2,4,6-triiodophthalamide (iohexol).

Similarly, in the production of iopentol as described in NO-160918, 5-Amine is acetylated to yield 5-Acetamide which is N-alkylated to produce 5-[N-(2-hydroxy-3-

methoxypropyl) acetamido-N,N'-bis(2,3-dihydroxypropyl) - 2,4,6-triiodophthalamide (iopentol).

In the production of iodixanol described in NO-161368, 5-Amine is acetylated to yield 5-Acetamide which is then reacted with a bifunctional alkylating agent, a coupling agent, to yield 1,3-bis(acetamido)-N,N'-bis[3,5-bis(2,3-dihydroxypropylaminocarbonyl)-2,4,6-triiodophenyl]-2-hydroxypropane (iodixanol).

In the preparation of iohexol described above, the acylation is effected in acetic anhydride in the presence of a catalytic amount of sulphuric acid followed by concentration and addition of methanol. Thereafter water and sodium hydroxide (to pH 10-11) are added, base hydrolysis (to remove O-acyl groups) proceeds for 4-5 hours and 5-Acetamide is isolated from the reaction mixture by cooling and neutralization with hydrochloric acid. The precipitated 5-Acetamide is filtered, washed with water and dried. The 5-Acetamide is then dissolved in propylene glycol with the addition of sodium methoxide, the resulting methanol is stripped off and the N-alkylating agent (1-chloro-2,3propanediol) is added. After the N-alkylation reaction is complete, the reaction mixture is evaporated to dryness and further purification steps (involving crystallization from a further solvent, butanol) are performed to yield the purified iohexol in a form suitable for use in secondary production.

The 5-Amine to 5-Acetamide acylation described for the preparation of iopentol in NO-160918 is similar, with the 5-Amine being acylated in acetic anhydride in the presence of a catalytic amount of p-toluene sulphonic acid. After cooling the reaction mixture, a precipitate forms which is filtered off and suspended in a mixture of methanol and water and hydrolysed under basic

conditions (pH 11.5). The 5-Acetamide product is filtered off after cooling to ambient temperature, neutralization with HCl and further cooling to 3°C. The 5-Acetamide is washed with water and dried before being suspended in propylene glycol. Sodium hydroxide is added and when all solid material has dissolved, the N-alkylating agent (here 1-chloro-3-methoxy-2-propanol) is added. After the alkylation is quenched, several purification steps are again required before satisfactorily pure iopentol is obtained.

For the synthesis of iodixanol described in NO-161368, 5-Acetamide is prepared and worked up as described for iopentol above. It is then suspended in water and dissolved therein with the addition of sodium hydroxide. The N-alkylating agent, the coupling reagent epichlorohydrin, is then added. After the reaction is complete it is quenched with dilute hydrochloric acid and further purification steps are carried out in order to obtain satisfactorily pure iodixanol.

#### Summary of the Invention

It has now been surprisingly found that the work-up of the 5-Acetamide before the subsequent N-alkylation may be avoided without unacceptable loss in yield or purity of the acylated, N-alkylated product and without undue complication of the purification procedure for that product.

Thus viewed from one aspect the invention provides a process for the preparation of an N-alkyl-acylamino-phenyl-carboxylic acid or carboxylic acid derivative by liquid phase acylation and subsequent N-alkylation of a corresponding amino-phenyl-carboxylic acid (or carboxylic acid derivative) characterised in that said N-alkylation is effected by addition of an alkylating

agent to a solution containing the reaction products of said acylation, ie. those that remain in solution during the liquid phase acylation.

Viewed from an alternative aspect the invention provides a process for the preparation of an N-alkyl-acylamino-phenyl-carboxylic acid or carboxylic acid derivative compound comprising acylating an amino-phenyl-carboxylic acid (or carboxylic acid derivative) in a liquid phase, base hydrolysing the acylated product to remove O-acyl groups from the N-acyl-amino intermediate and, maintaining the liquid phase at a basic pH, N-alkylating the N-acylamino intermediate.

## Detailed Description of the Invention

In the process of the invention, the aminophenyl carboxylic acid or carboxylic acid derivative starting product is preferably a compound having a total of three amino and carboxyl groups on the phenyl ring, eg. an aminoisophthalic acid or derivative or a diaminobenzoic acid or derivative. By carboxylic acid derivative is meant for example a salt, ester or amide, eg. an CONHR\* or COOR\* group where R\* is optionally hydroxylated alkyl preferably optionally hydroxylated  $C_{1-6}$  alkyl. Furthermore the starting compound is preferably a triiodophenyl compound, particularly an alkylaminocarbonyl-triiodophenyl compound and most particularly a 2,4,6-triiodo-2,5-bis(alkylamino-carbonyl)-aniline, such as 5-Amine for example. The alkyl moiety of any alkylaminocarbonyl group will preferably carry one or more hydroxyl groups and will typically contain up to 6, preferably up to 4 carbon atoms.

The acylation of the starting compound may be effected using any convenient acylating agent, eg. an acetylating agent such as for example an acid halide or more

preferably acetic anhydride.

Following the acylation, which is conveniently acid catalysed, the product is base hydrolysed, eg. at pH 11 to 12, to remove unwanted O-acyl groups without displacing the desired N-acyl groups. Base hydrolysis may be with any organic or inorganic base but is preferably effected using an inorganic base, eg. an alkali metal hydroxide such as sodium hydroxide.

After the base hydrolysis, the solution pH is if necessary adjusted to an appropriate basic value for N-alkylation (generally within the range 10-12), and if desired further solvent (eg. a lower alcohol such as a  $C_{1-4}$  alcohol, eg. methanol, optionally together with water) is added to the reaction mixture before, during or after addition to the reaction mixture of the alkylating agent, thereby ensuring that the non-isolated intermediate does not precipitate out.

Thus unlike prior successive acylation and N-alkylation reactions, the reaction medium is maintained at a basic pH and the N-acylated product (and the by-products) are not isolated from the reaction mixture before the subsequent N-alkylation step.

The N-alkylating agent used in the N-alkylation of the non-isolated intermediate may be any alkylating agent capable of introducing the desired alkyl group (or a precursor for a desired group) at the acylamino nitrogen. Where the acylated, N-alkylated product is intended to be a dimer (ie. containing two iodophenyl groups), the alkylating agent may be a bifunctional coupling agent.

For the preparation of iohexol, 1-halo-2,3-propanediols (eg. 1-chloro-2,3-propanediol) and glycidol are

preferred alkylating agents. For the preparation of iopentol, 1-halo-3-methoxy-2-propanols (eg. 1-chloro-3-methoxy-propanol) are preferred. For the production of iodixanol a coupling agent such as a 1,3-halo-2-propanol or more preferably epichlorohydrin may be used.

The alkylation reaction may be terminated conventionally by quenching with acid, eg. HCl, and the acetylated, N-alkylated product may also be worked up conventionally, eg. by crystallization from an appropriate solvent or solvent mixture, eg. one or more  $C_{1-6}$  alcohols.

Where the N-alkyl-acylamino product is a product such as iohexol, iopentol or iodixanol, this work-up yields a product which can be used for secondary production of an X-ray contrast medium. However for other end products, the product may be used as the starting material for further synthetic steps, eg. in the primary production of a non-ionic X-ray contrast agent.

Thus the process of the present invention allows N-acylation and N-alkylation to be effected as a one-pot synthesis achieving savings in equipment, time and materials without undue loss in yield and, surprisingly, achieving in the purification of the end product comparable or better purity levels to those achieved when the intermediate and the end product are both subject to work up stages.

The process of the present invention is particularly applicable to the preparation of the following N-alkyl-acylaminophenyl X-ray contrast agents: iomeprol, ioversol, ioxilan, iotrolan, ioxaglate, iodecimol, 2-iopyrol, 2-iopiperidol, iohexol, iopentol and iodixanol.

The invention will now be illustrated further by reference to the following non-limiting Examples.

#### EXAMPLE 1

5-Amino-N,N'-bis(2,3-dihydroxypropyl)-2,4,6-triiodoisophthalamide (388 g), acetic anhydride (838 g) and p-toluene sulfonic acid (2 g) were mixed and heated to reflux for 1.5 hours, before concentration to a highly viscous solution under reduced pressure.

Methanol (44 g) was added to the solution in several portions with distillation under atmospheric pressure between each addition, followed by a final portion of methanol (266 g). A small amount of distilled water (40 ml) was added before stirring at 55°C overnight.

A portion (490 ml) of a solution made as described above was added to a jacketed reactor at about 30°C, and stirred with a speed of 150 rpm. The pH was raised by aqueous sodium hydroxide (50 w/v%, 190 ml) to about 12. 1-Chloro-2,3-propanediol (49 g) was added to the solution, and the pH was further adjusted by sodium hydroxide (50 w/v%, 25 ml) to about 11.5. Small amounts of additional 1-chloro-2,3-propanediol (total 6 ml) were added during the first 24 hours of the reaction. After a total reaction time of 99 hours, the solution was analyzed by reversed phase HPLC (water/acetonitrile as eluent) with the following results:

Iohexol	94.4%
5-Acetamide	1.9%
5-Amine	0.6%
Other impurities	3.1%

#### EXAMPLE 2

5-Amino-N,N'-bis(2,3-dihydroxypropyl)-2,4,6triiodoisophthalamide (388 g), acetic anhydride (838 g) and p-toluene sulfonic acid (2 g) were mixed and heated to reflux for 1.5 hours, before concentration to a highly viscous solution under reduced pressure.

Methanol (44 g) was added to the solution in several portions with distillation under atmospheric pressure between each addition, followed by a final portion of methanol (266 g). A small amount of distilled water (40 ml) was added before stirring at 55°C overnight.

A portion (490 ml) of a solution made as described above was added to a jacketed reactor at about 35°C, and stirred with a speed of 150 rpm. The pH was raised by aqueous sodium hydroxide (50 w/v%, 200 ml) to about 12. Additional sodium hydroxide solution (10 ml) and 1-chloro-3-methoxy-2-propanol (42 g) were added, and the pH was adjusted back to about 12 by small amounts of sodium hydroxide solution. Further additions of 1-chloro-3-methoxy-2-propanol (total 21 g) were added after 2, 3, 4, 5 and 6 hours of reaction. After a total reaction time of 53 hours, the mixture was analyzed by HPLC (water/acetonitrile as eluent) with the following results:

Iopentol	91.4%
5-Acetamide	1.6%
5-Amine	0.3%
Other impurities	6.8%

#### Claims:

- 1. In a process for the preparation of an N-alkyl-acylamino-phenyl-carboxylic acid or carboxylic acid derivative by liquid phase acylation and subsequent N-alkylation of a corresponding amino-phenyl-carboxylic acid (or carboxylic acid derivative), the improvement comprising the addition of an alkylating agent to a solution containing the reaction products of said acylation, to effect said N-alkylation.
- 2. A process as claimed in claim 1 wherein said corresponding amino-phenyl carboxylic acid (or carboxylic acid derivative) is a compound having a total of three amino and carboxyl groups on the phenyl ring thereof.
- 3. A process as claimed in claim 1, wherein said corresponding amino-phenyl-carboxylic acid (or carboxylic acid derivative) is a triiodophenyl compound.
- 4. A process as claimed in claim 1, wherein said corresponding amino-phenyl-carboxylic acid (or carboxylic acid derivative) is an alkylamino-carbonyl-triiodophenyl compound.
- 5. A process as claimed in claim 1, wherein said corresponding amino-phenyl-carboxylic acid (or carboxylic acid derivative) is a 2,4,6-triiodo-2,5-bis(alkylamino-carbonyl)-aniline.
- 6. A process as claimed in claim 1, wherein said corresponding amino-phenyl-carboxylic acid (or carboxylic acid derivative) is 5-amino-N-N'-bis(2,3-dihydroxypropyl)-2,4,6-triiodophthalamide.

- 7. A process as claimed in claim 1, wherein said corresponding amino-phenyl-carboxylic acid (or carboxylic acid derivative) contains an alkylaminocarbonyl group carrying one or more hydroxyl groups and containing up to 6 carbon atoms.
- 8. A process as claimed in claim 1, wherein said acylation is effected using an acid halide or acetic anhydride.
- 9. A process as claimed in claim 1, wherein said alkylating agent is selected from the group consisting of 1-halo-2,3-propanediols, glycidol, 1-halo-3-methoxy-2-propanols, 1,3-halo-2-propanols and epichlorohydrin.
- 10. A process for the preparation of an N-alkyl-acylamino-phenyl-carboxylic acid or carboxylic acid derivative compound comprising acylating an amino-phenyl-carboxylic acid (or carboxylic acid derivative) in a liquid phase, base hydrolysing the acylated product to remove O-acyl groups from the resulting N-acyl-amino intermediate and, maintaining the liquid phase at a basic pH, N-alkylating said N-acylamino intermediate.
- 11. A process as claimed in claim 1 or claim 10, being a process for the preparation of a contrast agent selected from the group consisting of iomeprol, ioversol, ioxilan, iotrolan, ioxaglate, iodecimol, 2-iopyrol, 2-iopiperidol, iohexol, iopentol and iodixanol.

#### Summary

# Process for the preparation of contrast agents

The invention provides a process for the preparation of an N-alkyl-acylamino-phenyl-carboxylic acid or carboxylic acid derivative by liquid phase acylation and subsequent N-alkylation of a corresponding amino-phenyl-carboxylic acid (or carboxylic acid derivative), the improvement comprising the addition of an alkylating agent to a solution containing the reaction products of said acylation, to effect said N-alkylation.

ATTORNEY/DOCKET NO: REF/Dehn

# DECLARATION FOR PATENT APPLICATION AND APPOINTMENT OF ATTORNEY

As a below named inventor, I hereby declare that my residence, post office address and citizenship are as stated below next to my name; I believe that I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention (Design, if applicable) entitled:

PROCESS FOR THE PREPARATION OF CONTRAST ACENTS.

PROCESS FOR THE PREPARA	TION OF CONTRAST AGENTS	S		11	,
the specification of which (check o	ne):				
X is attached hereto, or ☐ was filed on:  Number:		as U.	S. Application Number or PCT	Internation	al Application
and (if applicable) was	amended on:				
I hereby state that I have reviewed amendment(s) referred to above. I of Federal Regulations, §1.56. I h for patent or inventor's certificate I a filing date before that of the appl	acknowledge the duty to disclose in tereby claim foreign priority benefit isted below and have also identified	nformation w ts under <i>Title</i> l below anv f	hich is material to patentability as	defined in	Title 37, Cod
	PRIOR FOREIGN APPLICATI	on(s)		PRIORITY	CLAIMED
Number	Country		Day/Month/Year Filed	Yes	No
96 18055.9	Great Britain		29 August 1996	X	
	☐ Additional Priority Application	on(s) Listed of	on Following Page(s)		
	T UNDER TITLE 35 U.S. CODE §1	19(E) OF AN	Y U.S. PROVISIONAL APPLICATION	NS LISTED B	ELOW.
	on Number		Day/Month/Year Fil	ed	
5 2 3 3 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4	60/029,143	3		21 O	ctober 1996
9 WA .	☐ Additional Provisional Applica	tion(s) Listed	on Following Page(s)		
Application Number	application(s) and the national or P	C1 Internation	Status - Patented, Pendi		adonad
- 102 H			Otatus - Fatenteu, Fendi	ing of Abai	ndoned
	Additional US/PCT Priority Appl	ication(s) list	ed on Following Page(s)		
punishable by fine or imprisonment, eopardize the validity of the application POWER OF ATTORNEY: I his application and transact all businger, Reg. No. 25,893; Richard E.	or both, under section 1001 of title ation or any patent issued thereon.  (We) hereby appoint as my (our) ness in the Patent and Trademark O Fichter, Reg. No. 26.382; Charles	ith the knowled 18 of the Ur attorneys, wiffice connect	edge that willful false statements wited States Code and that such wi th full powers of substitution and ed therewith: J. Ernest Kenney, I	and the like llful false st revocation	e so made are atements may  to prosecute
oruce H. 170xell, Reg. No. 26,392	; and accept and follow instructions from				, ,
egarding any matter related to the p	preparation, examination, grant and	l maintenance	e of this application, any continua	tion, contin	uation-in-part
end correspondence to: BACO 625 Sla	y patent resulting therefrom, until I( N & THOMAS aters Lane - 4th Floor dria, VA 22314		ur) assigns withdraw this authoriz one Calls to: Richard E (703) 683-0500		iting.
FULL NAME OF FIRST OR SOLE INVENT Trygve GULBRANDSEN	OR	CITIZENSHIP Nors	vegian		
RESIDENCE ADDRESS			ADDRESS IS THE SAME AS RESIDENCE	E ADDRESS	UNLESS
c/o Nycomed Imaging AS Nycoveien 1-2, N-0401 Osl	o 4, Norway		SHOWN BELOW		
DATE Am 1 10, 199	2	SIGNATURE	~ O 11 1		

X See following page(s) for additional joint inventors.

ATTORNEY/DOCKET NO: REF/Dehn

# CONTINUATION OF DECLARATION FOR PATENT APPLICATION AND APPOINTMENT OF ATTORNEY

Page \_\_2\_

II = N	
FULL NAME OF JOINT INVENTOR	CITIZENSHIP
Odd Einar INGVOLDSTAD	Norwegian
RESIDENCE ADDRESS	POST OFFICE ADDRESS IS THE SAME AS RESIDENCE ADDRESS UNLESS
c/o Nycomed Imaging AS	OTHERWISE SHOWN BELOW
Nycoveien 1-2, N-0401 Oslo 4, Norway	
DATE COCK COCK TO COCK	SIGNATURE CO. 1. A
08.04.1997	Odd E. Inguldstad
F N	
FULL NAME OF JOINT INVENTOR Lars Torio HOLMAAS	CITIZENSHIP
Lars Terje HOLMAAS	Norwegian
RESIDENCE ADDRESS	POST OFFICE ADDRESS IS THE SAME AS RESIDENCE ADDRESS UNLESS
c/o Nycomed Imaging AS	OTHERWISE SHOWN BELOW
Nycoveien 1-2, N-0401 Oslo 4, Norway	
DATE 4/4 07	SIGNATURE
DATE 4/4-97	Jano 198 Holman
FULL NAME OF JOINT INVENTOR	CITIZENSHIP
700ats. 2 12 7-mars	
RESIDENCE ADDRESS	POST OFFICE ADDRESS IS THE SAME AS RESIDENCE ADDRESS UNLESS
STORMS  TO SEE SEE SEE SEE SEE SEE SEE SEE SEE SE	OTHERWISE SHOWN BELOW
And The State of t	
CONTROL OF THE CONTRO	
DATE	SIGNATURE
	GIGINATORE
L I I	
FULL NAME OF JOINT INVENTOR	CITIZENSHIP
E SOURCE OF SOURT HANDERSON	CITIZENSHIP
RESIDENCE ADDRESS	
HESIDENCE ADDRESS	POST OFFICE ADDRESS IS THE SAME AS RESIDENCE ADDRESS UNLESS
	OTHERWISE SHOWN BELOW
i i i i i i i i i i i i i i i i i i i	
2	
DATE	SIGNATURE

PATAPATT.DEC (081596)

 $<sup>\</sup>hfill \square$  See following pages for additional joint inventors/priority applications.

## IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of:

Trygve GULBRANDSEN et al.

Attention:

Filed: Herewith on April 21, 1997

**Applications Branch** 

For:

PROCESS FOR THE PREPARATION

OF CONTRAST AGENTS

#### PRELIMINARY AMENDMENT

Assistant Commissioner for Patents Washington, D.C. 20231

Sir:

Prior to an examination on the merits, please amend the above identified application as follows:

#### IN THE CLAIMS:

Claim 11, line 1, delete "or claim 10".

Please add the following claim to the application:

--12. A process as claimed in claim 10, being a process for the preparation of a contrast agent selected from the group consisting of iomeprol, ioversol, ioxilan, iotrolan, ioxaglate, iodecimol, 2-iopyrol, 2-iopiperidol, iohexol, iopentol and iodixanol.--

#### **REMARKS**

Applicants have amended the claims in order to remove the multiple dependency from claim 11 and reintroduce the cancelled subject matter as new claim 12.

Respectfully submitted,

Richard E. Fichter

Registration No. 26,382

BACON & THOMAS 625 Slaters Lane - 4th Fl. Alexandria, Virginia 22314 Phone: (703) 683-0500

REF:TJL:ti

April 21, 1997